

Information provided is intended to inform clients of expected procedures and outcomes. However, not all patients, procedures or surgeons are the same, and information may not be applicable to all procedures. PIAC reserves the right to vary the way that a procedure is performed or which technique is used.

Breast Reduction

PRICE (THB)
140,000

Surgical/Procedure Name

Reduction Mammoplasty

Common Name

Breast Reduction

Introduction

In recent years breast reduction has become a very common procedure. Very large and heavy breasts can lead to a variety of medical problems, such as back and neck pain, skin irritation, skeletal deformities and breathing problems. As well as medical problems, unusually large breasts can also make women feel extremely self-conscious. The aim of the procedure is to produce smaller, shapelier breasts, which are in proportion with the rest of the body. The size of the areola, the darker skin surrounding the nipple, can also be reduced.

Surgical Procedure

During surgery, breast fat, glandular tissue, and skin are removed, to make the breasts smaller, lighter, and firmer. An incision is usually made around the nipple and under the surface of the breast, like an upside down T. The nipple and areola are almost always left attached to their blood vessels and nerves. The skin from above the nipples is then pulled down to shape the new breast and a hole is cut for the new nipple and areola. If the breasts are particularly large the nipple may need to be completely removed and grafted to a higher position - this results in a loss of sensation to the nipple and areola.

Hospital Admission

Two nights

Duration of Operation

Four to six hours


Anaesthetic

General anaesthesia

Pre Operative Care

Before surgery please inform your surgeon of any allergies, all medical conditions, and any medication that you are taking (both prescription and non-prescription). To eliminate the chance of post op. bleeding you should avoid aspirin and any medication containing aspirin for two weeks prior to surgery. You should also not smoke for 2 weeks prior to surgery as smoking can affect your reaction to the anaesthetic and slow down the healing process. Patients that suffer from hypertension must inform the surgeon prior to surgery.

In your consultation you should discuss your expectations of the outcome of the surgery and listen to the surgeon's opinion. You will probably be required to send photos of your breasts to the surgeon prior to the consultation. This will enable the assessment and also serve as a reference following surgery.



Post Operative Care

When you wake after surgery you will be wrapped in elastic bandages, gauze dressings and a surgical bra. A small tube may be placed in each breast to drain blood and fluids for the first day or two.

The extent of the post operative swelling and bruising depends on whether you tend to bruise or swell easily. Every person is different. Application of cold compresses or ice packs will reduce swelling and relieve discomfort.

A couple of days after surgery the pressure bandages will be removed. The surgical bra must be worn at all times for several weeks, until the swelling and bruising subsides, or until your surgeon advises you otherwise. You should only remove the surgical bra to bathe and to wash the garment.

Usually, sutures will be removed in the first week. Sometimes the surgeon will use dissolvable sutures. If the breast skin is very dry following surgery, a moisturizer can be applied several times a day. Vitamin E cream is extremely effective and also helps the scarring to heal, however the suture area must be kept dry at all times, particularly after bathing.

For the first few months it is important not to place tension on the incisions by performing strenuous activities like lifting or stretching your arms above your head, as this may increase scarring by stretching the tissue. You will need assistance to carry your groceries or hang your washing !

Your surgeon will try to ensure that your incisions are as subtle as possible, however it is important to remember that the scars are extensive and permanent. They may be red for months, and then slowly become fainter, eventually fading to thin white lines. In most cases the scars are positioned so that they would not be seen when wearing low-cut tops and dresses.

Recovery usually takes about two weeks. Most people are back to work within three weeks.

Risks and Complications

Risks are inherent to any surgical procedure. The most common risks are swelling, bruising, bleeding, infection, fluid, scarring, numbness, or a change in sensation to the nipple. The most common risk particular to this surgery is damage to the nerve and blood supply of the nipple. The possibility of breast feeding may decrease after surgery. Breast reduction is recommended for people that have had children already and do not wish to have any more. Breast reduction does NOT increase the risk of breast cancer and indeed may slightly decrease the risk. Mammograms will still be possible after surgery.

Scars will fade gradually but nipple numbness may take up to 12 months

